

# Single spherical particle settling: a benchmarking data-set generated by spectral/spectral-element DNS

A full description of the simulations is available in reference [4].

## Description of the flow

A single spherical particle (rigid, with homogeneous mass density) is freely settling under the influence of gravity in an ambient, viscous, incompressible fluid.

## Geometry

The computational domain is cylindrical, with the cylinder axis aligned with the direction of gravity. The particle is located on the axis of the cylinder, at a distance  $L_u$  from the inflow and at  $L_d$  from the outflow plane (cf. sketch in figure 1 and table 1).

## Flow parameters

The problem is governed by two parameters. The first is the solid-to-fluid density ratio  $\rho_p/\rho_f$  which is maintained constant at a value of 1.5 in this study. The second is the Galileo number  $G = u_g d/\nu$ , where  $u_g = (|\rho_p/\rho_f - 1| d |\mathbf{g}|)^{1/2}$  is the gravitational velocity scale,  $d$  is the particle diameter,  $\mathbf{g}$  the vector of gravitational acceleration and  $\nu$  the kinematic viscosity of the fluid. The Galileo number was varied in the range from 144 to 250 (cf. table 2), leading to four different flow regimes and patterns of particle motion.

- Case “A”: axi-symmetric flow, steady vertical particle path;
- case “B”: oblique double-threaded wake, steady oblique path;
- case “C”: periodic hairpin vortex shedding, oscillating oblique path;




Figure 1: The geometry of the problem and the computational domain.




Figure 2: Sketch of the notation concerning directional unit vectors and coordinates in the plane spanned by the vertical axis  $\mathbf{e}_z$  and the direction of the particle motion  $\mathbf{e}_{p\parallel}$ , as defined in [4]. Note that the direction  $\mathbf{e}_{pH z\perp}$  is perpendicular to the plane of the sketch which corresponds to the plane defined by the trajectory and the vertical.

$D_\Omega$	$L_u$	$L_d$	case	A	B	C	D
7.54	5	10	$Ga$	144	178.46	190	250

Table 1: Dimensions of the computational domain: horizontal diameter  $D_\Omega$  and distance from the sphere’s center to the up-/downstream boundaries  $L_u$  and  $L_d$ .

Table 2: Values of the Galileo number  $G$  chosen in each one of the 4 flow cases. In all cases the solid-to-fluid density ratio was set to  $\rho_p/\rho_f = 1.5$ .

- case “D”: chaotic wake and particle path.

## Numerical method and resolution

The method solves the incompressible Navier-Stokes equations on a mesh which is translating with the sphere, thereby avoiding remeshing. The spatial discretization uses a truncated Fourier series in the azimuthal direction and a spectral-element approach in the radial/vertical plane. The temporal discretization is semi-implicit using a third order Adams-Bashforth method for the non-linear terms. Momentum and continuity equations are uncoupled through a projection method. Details can be found in [1; 2].

- Grid: 134 elements ( $G = 144$  and  $G = 178$ ), 169 elements ( $G = 190$  and  $G = 250$ ).
- 6 collocation points in each of the two spatial directions internal to each element.
- Truncation of azimuthal Fourier series at wave-number 7.
- Time step: such that  $CFL \approx 0.25$ .

## Boundary conditions

- inflow: zero (ambient) velocity.
- side-walls and outflow: no stress Neumann condition
- pressure: homogeneous Dirichlet condition on all Neumann boundaries of velocity.

## Available data

The fluid velocity data is presented relative to the particle velocity. A Cartesian coordinate system attached to the particle center is used which is constructed as follows. The “local trajectory plane” (shown in figure 2) is defined by  $\mathbf{g}$  and by the particle velocity vector; the latter one defines the unit vector  $\mathbf{e}_{p\parallel}$ . The vector oriented normal to that plane is denoted by  $\mathbf{e}_{pHz\perp}$ . The two unit vectors  $\mathbf{e}_{p\parallel}$  and  $\mathbf{e}_{pHz\perp}$  are completed by  $\mathbf{e}_{p\perp}$  which lies in the trajectory plane and which is perpendicular to the particle trajectory.

The data-set contains the following data items:

- relative velocity deficit on the axis through the particle;
- cross-profiles of the three (relative) velocity components at 4 distances downstream of the particle, taken along the directions  $\mathbf{e}_{pHz\perp}$  and  $\mathbf{e}_{p\perp}$ ;
- pressure coefficient on two great circles (in the trajectory plane and in the perpendicular vertical plane);
- temporal evolution of the different particle degrees of freedom (time-periodic case “C” only);
- probability density function and temporal auto-correlation of the particle degrees of freedom (chaotic case “D” only).

Please note that all lengths given herein and in the data-base are **normalized with the particle diameter  $d$** ; all velocity data is **normalized with the gravitational velocity  $u_g$** .

## Data format and location

Data is presented in the form of “mat-files”<sup>\*</sup> (readable e.g. with Matlab and GNU/octave). A script for reading the data (and plotting it) is provided. The data is located below the following URL:

[www.ifh.kit.edu/dns\\_data/particles/single\\_sphere\\_sedimentation](http://www.ifh.kit.edu/dns_data/particles/single_sphere_sedimentation)

## Contact

Markus Uhlmann

Institute for Hydromechanics

Karlsruhe Institute of Technology (KIT)

76131 Karlsruhe, Germany

[markus.uhlmann@kit.edu](mailto:markus.uhlmann@kit.edu)

Jan Dušek

Institut de Mécanique des Fluides et des Solides

Université de Strasbourg

67000 Strasbourg, France

[dusek@unistra.fr](mailto:dusek@unistra.fr)

## References

- [1] B. Ghidersa and J. Dušek. Breaking of axisymmetry and onset of unsteadiness in the wake of a sphere. *J. Fluid Mech.*, 423:33–69, 11 2000. [doi:10.1017/S0022112000001701](https://doi.org/10.1017/S0022112000001701).
- [2] M. Jenny and J. Dušek. Efficient numerical method for the direct numerical simulation of the flow past a single light moving spherical body in transitional regimes. *J. Comput. Phys.*, 194:215–232, 2004.
- [3] M. Jenny, J. Dušek, and G. Bouchet. Instabilities and transition of a sphere falling or ascending freely in a Newtonian fluid. *J. Fluid Mech.*, 508:201–239, 2004.
- [4] M. Uhlmann and J. Dušek. The motion of a single heavy sphere in ambient fluid: a benchmark for interface-resolved particulate flow simulations with significant relative velocities. *Int. J. Multiphase Flow*, 2013. (*accepted for publication*). [arXiv:1310.5358](https://arxiv.org/abs/1310.5358), [doi:10.1016/j.ijmultiphaseflow.2013.10.010](https://doi.org/10.1016/j.ijmultiphaseflow.2013.10.010).

---

<sup>\*</sup>A detailed description of the file format can be found here: [http://www.mathworks.com/help/pdf\\_doc/matlab/matfile\\_format.pdf](http://www.mathworks.com/help/pdf_doc/matlab/matfile_format.pdf)